Open Horizon Scientific Review

Vol. 02 No.02 2024

Biochemical Pathways Involved in the Decomposition of Complex Organic Substances

Ayesha Noreen[1]

[1] Institute of Biochemistry and Biotechnology, University of Agriculture, Faisalabad, Pakistan

ABSTRACT

The decomposition of complex organic substances such as cellulose, hemicellulose, lignin, proteins, lipids and aromatic compounds is a central process in the global carbon and nitrogen cycles. It is carried out mainly by bacteria, fungi and archaea that secrete specialized enzymes and operate interconnected catabolic pathways. This article reviews the biochemical routes through which large polymeric molecules are depolymerized, converted into small intermediates and finally oxidized to carbon dioxide or reduced to methane. The roles of extracellular hydrolases and oxidoreductases, the degradation of lignin by peroxidases and laccases, proteolysis and nitrogen mineralization, lipid beta-oxidation, aromatic ring cleavage and anaerobic fermentation are described in detail. Environmental controls such as temperature, moisture, oxygen availability and substrate chemistry are discussed, together with applications in composting, biofuel production and waste management. Understanding these pathways at molecular level provides a basis for improving bioremediation, enhancing soil fertility and developing efficient biorefinery processes.

Key words : *biochemical pathways, organic matter decomposition, lignocellulose, microbial enzymes, anaerobic digestion*

Introduction:

Every year photosynthesis fixes an enormous quantity of carbon into plant biomass, and almost all of this material is eventually returned to the atmosphere or to the soil through decomposition. Without this recycling, nutrients would remain locked in dead tissue, and the productivity of terrestrial and aquatic ecosystems would collapse. Decomposition is therefore not simply a process of decay but a fundamental metabolic service that links primary production with nutrient availability, soil formation and greenhouse gas fluxes.The organic substances that must be broken down are chemically diverse. Plant cell walls consist of crystalline cellulose embedded in a matrix of hemicellulose and the irregular aromatic polymer lignin. Animal and microbial remains contribute proteins, nucleic acids, lipids and chitin, while human activities add synthetic and semi-synthetic organic compounds, many of which resemble natural aromatic molecules. Because these substances differ greatly in bond type, solubility and molecular size, no single enzyme or organism can degrade them all. Instead, decomposition proceeds through sequential and cooperative pathways in which the products of one organism become the substrates of another.In the first stage, large polymers that cannot cross the cell membrane are attacked outside the cell by secreted hydrolytic and oxidative enzymes. In the second stage, the resulting monomers and small fragments are taken up and funneled into central metabolic routes such as glycolysis, the

pentose phosphate pathway and the tricarboxylic acid cycle. In the final stage, electrons are transferred to oxygen or to alternative acceptors in respiration, or are released as reduced end products in fermentation and methanogenesis. The nature of the available electron acceptor determines how much energy is gained and which end products appear.This article provides an organized account of these pathways. It begins with the composition of organic matter and the microbial communities involved, then describes the principal enzymatic routes for polysaccharides, lignin, proteins, lipids and aromatic compounds. It continues with anaerobic processes and closes with the environmental factors, applications and research directions that are most relevant for agriculture, environmental protection and biotechnology, followed by a brief summary.

Overview of Organic Matter Composition and Decomposition:

Organic matter in terrestrial ecosystems is dominated by plant-derived polymers. Cellulose, a linear chain of beta-1,4-linked glucose units that aggregate into microfibrils, typically makes up roughly one third to one half of the dry mass of plant tissue. Hemicelluloses are branched heteropolymers of xylose, mannose, arabinose and other sugars, and they cross-link with cellulose and lignin. Lignin, formed by radical coupling of phenylpropanoid alcohols, is an amorphous and heterogeneous network that gives rigidity to wood and protects polysaccharides from enzymatic attack. Proteins, lipids, starch and soluble sugars are present in smaller amounts but are decomposed much faster and supply most of the readily available energy and nitrogen.Decomposition is usually described as a sequence of three overlapping processes. Leaching removes soluble compounds shortly after tissue death, fragmentation by soil fauna increases the surface area available to microorganisms, and biochemical catabolism converts the organic carbon to carbon dioxide, biomass and humified residues. The rate at which these processes proceed depends on the chemical quality of the substrate, with high lignin content and high carbon-to-nitrogen ratio generally slowing decomposition, and on the physical environment. The graph below compares the approximate extent of mass loss of major components of plant litter after a short incubation, which illustrates the strong dependence of decomposition rate on molecular structure.

Microbial Communities and Enzymatic Machinery:

Decomposition is carried out by a succession of microbial groups. Bacteria, especially those belonging to the Actinobacteria, Proteobacteria and Firmicutes, dominate the early utilization of soluble compounds and proteins, whereas fungi, particularly the white-rot and brown-rot basidiomycetes and many ascomycetes, are the most effective degraders of lignocellulose in aerated environments. In oxygen-free habitats, consortia of fermentative bacteria, syntrophic acetogens and methanogenic archaea take over, forming metabolic chains in which hydrogen and acetate are rapidly consumed to keep the thermodynamics of each step favorable.The enzymatic toolkit of these organisms is extremely diverse. The Carbohydrate-Active Enzymes classification groups glycoside hydrolases, polysaccharide lyases, carbohydrate esterases and auxiliary activities into families based on sequence similarity, and it shows that cellulolytic organisms often encode dozens of related enzymes with complementary specificities. Proteases,

lipases, phosphatases and nucleases supply the other nutrient-releasing activities. Oxidative enzymes such as lignin peroxidase, manganese peroxidase, laccase and lytic polysaccharide monooxygenase act on substrates that are too recalcitrant for hydrolysis alone.Enzyme production is tightly regulated. Many degradative genes are induced by the substrate or by its breakdown products and repressed by easily metabolized sugars through catabolite repression. This regulation conserves cellular resources and also explains why the addition of labile carbon can either stimulate or suppress the breakdown of more complex material, a phenomenon known as priming. Enzyme activities measured in soils and composts are therefore widely used as indicators of the functional state of the microbial community.

Extracellular Hydrolysis of Polysaccharides:

Cellulose degradation is accomplished by the cooperative action of three major groups of hydrolytic enzymes. Endoglucanases cleave internal beta-1,4 bonds in amorphous regions, creating new chain ends, exoglucanases or cellobiohydrolases then remove cellobiose units processively from the chain ends, and beta-glucosidases hydrolyze cellobiose to glucose, relieving product inhibition. In many anaerobic bacteria these enzymes are assembled on a scaffold protein to form a large extracellular complex called the cellulosome, which brings the catalytic modules into close proximity with the crystalline substrate and with the cell surface.Aerobic fungi complement hydrolysis with lytic polysaccharide monooxygenases, copper-containing enzymes that oxidatively cleave glycosidic bonds on the surface of crystalline cellulose and thereby create access points for hydrolases. The electrons required for this reaction are provided by small reductants such as phenolic compounds derived from lignin or by cellobiose dehydrogenase. This discovery has changed the traditional view of cellulose degradation as a purely hydrolytic process and has had direct consequences for the design of industrial enzyme cocktails.Hemicellulose degradation requires a wider range of enzymes because of its heterogeneous structure. Endo-xylanases cut the xylan backbone, beta-xylosidases release xylose, and accessory enzymes such as arabinofuranosidases, acetyl xylan esterases and feruloyl esterases remove side chains and ester links to lignin. Starch is attacked by alpha-amylases and glucoamylases, pectin by polygalacturonases and pectate lyases, and chitin by chitinases and N-acetylglucosaminidases. The resulting hexoses and pentoses enter glycolysis or the pentose phosphate pathway after uptake by specific transporters.

Lignin Degradation Pathways:

Lignin cannot be hydrolyzed because its subunits are connected by stable carbon-carbon and ether bonds rather than by regular hydrolyzable linkages. Its breakdown therefore depends on oxidative reactions that generate reactive radicals. White-rot fungi secrete lignin peroxidase, manganese peroxidase and versatile peroxidase, which use hydrogen peroxide as oxidant. Lignin peroxidase oxidizes non-phenolic aromatic rings directly, while manganese peroxidase oxidizes Mn(II) to Mn(III), which is stabilized by organic acids such as oxalate and diffuses into the wood to oxidize phenolic structures.Laccases are multicopper oxidases that reduce molecular oxygen to water while oxidizing phenolic substrates, and in the presence of small redox mediators they

can also attack non-phenolic units. Hydrogen peroxide is supplied by enzymes such as glyoxal oxidase and aryl alcohol oxidase, which couple the oxidation of small aldehydes and alcohols to peroxide formation. The oxidation of the polymer produces unstable radicals that undergo side-chain cleavage, demethoxylation and aromatic ring opening, releasing smaller aromatic fragments.Bacterial lignin degradation is less well characterized but is increasingly recognized. Actinomycetes, Pseudomonas and Rhodococcus strains produce extracellular peroxidases, dye-decolorizing peroxidases and laccase-like enzymes, and they possess intracellular pathways such as the beta-aryl ether cleavage route that convert dimeric lignin fragments to vanillate, protocatechuate and related compounds. These intermediates are then processed by the aromatic catabolic pathways described below. The slow and incomplete oxidation of lignin also contributes to the formation of stable humic substances, which are important for soil carbon storage.

Proteolysis and Nitrogen Mineralization:

Proteins and peptides are the largest pool of organic nitrogen in most soils and decaying tissues. Extracellular proteases, including serine, metallo, cysteine and aspartic proteases, hydrolyze peptide bonds and release oligopeptides and amino acids. Because small peptides and free amino acids can be taken up directly by many microorganisms and by plant roots, the proteolytic step is considered the rate-limiting step in the supply of bioavailable nitrogen rather than the later conversion to ammonium.Inside the cell, amino acids are either incorporated into new proteins or deaminated to provide carbon skeletons and energy. Transaminases transfer amino groups to alpha-ketoglutarate to form glutamate, and glutamate dehydrogenase releases ammonium by oxidative deamination. The carbon skeletons enter the tricarboxylic acid cycle at the level of pyruvate, acetyl coenzyme A, oxaloacetate, alpha-ketoglutarate, succinyl coenzyme A or fumarate, depending on the amino acid. In anaerobic habitats, amino acids may be fermented by the Stickland reaction, in which one amino acid serves as electron donor and another as acceptor.Whether the released ammonium is retained or released to the environment depends on the nitrogen demand of the microbial community. When the substrate has a high carbon-to-nitrogen ratio, microorganisms immobilize ammonium into biomass, but when nitrogen is in excess it is mineralized and becomes available for nitrification, in which ammonia-oxidizing and nitrite-oxidizing microorganisms convert it first to nitrite and then to nitrate. Nucleic acids follow a similar course, with nucleases, phosphatases and deaminases releasing phosphate, ammonium and sugars.

Lipid Breakdown and Beta-Oxidation:

Lipids, including triacylglycerols, phospholipids and waxes, are energy-rich compounds that are degraded by lipases and phospholipases secreted by bacteria and fungi. Lipases hydrolyze the ester bonds of triacylglycerols at the lipid-water interface to release glycerol and long-chain fatty acids. Glycerol is phosphorylated and oxidized to dihydroxyacetone phosphate, which joins glycolysis, while fatty acids are activated to acyl coenzyme A derivatives and enter the beta-oxidation spiral.In each turn of beta-oxidation, the acyl chain is dehydrogenated, hydrated, dehydrogenated again and finally cleaved by thiolase, releasing one molecule of acetyl coenzyme A and a

chain shortened by two carbon atoms. The reaction also produces reduced flavin adenine dinucleotide and reduced nicotinamide adenine dinucleotide, whose electrons feed the respiratory chain. The acetyl coenzyme A is oxidized in the tricarboxylic acid cycle or, in organisms growing exclusively on fatty acids, passed through the glyoxylate cycle to generate four-carbon intermediates for gluconeogenesis.Unsaturated and odd-chain fatty acids require auxiliary enzymes such as isomerases, reductases and propionyl coenzyme A carboxylase. Under anaerobic conditions, syntrophic bacteria oxidize fatty acids to acetate and hydrogen, but this reaction is thermodynamically possible only when hydrogen is kept at very low concentration by methanogens or sulfate reducers. Accumulation of long-chain fatty acids may therefore inhibit anaerobic digestion in lipid-rich wastes, which is a well-known operational problem in biogas plants.

Aromatic Compound Catabolism:

Aromatic compounds originate from lignin, tannins, flavonoids, aromatic amino acids and anthropogenic pollutants, and their stable ring structure makes them difficult to degrade. Aerobic bacteria and fungi use a strategy in which a variety of peripheral pathways convert different substrates to a small number of central dihydroxylated intermediates such as catechol, protocatechuate and gentisate. Monooxygenases and dioxygenases introduce hydroxyl groups using molecular oxygen, which is both the co-substrate and the means of activating the otherwise unreactive ring.The central intermediates are cleaved by ring-fission dioxygenases. Intradiol dioxygenases cleave the bond between the two hydroxylated carbons to form muconic acid derivatives, whereas extradiol dioxygenases cleave adjacent to the hydroxyl groups to give semialdehyde products. In the beta-ketoadipate pathway, the products are converted through a series of reactions to succinyl coenzyme A and acetyl coenzyme A, which enter the tricarboxylic acid cycle. This pathway is widely distributed among soil bacteria and fungi and is a key route for the complete mineralization of lignin-derived monomers.In the absence of oxygen, an entirely different strategy is used. Anaerobic bacteria activate aromatic substrates by attaching coenzyme A and then reduce the ring by ATP-dependent or ATP-independent benzoyl coenzyme A reductases, after which the dearomatized ring is opened by hydrolysis and degraded by a modified beta-oxidation. This process allows the decomposition of aromatic compounds in sediments, flooded soils and digesters, although it is considerably slower than aerobic degradation and yields much less energy per mole of substrate.


AnaerobicDecomposition, Fermentation and Methanogenesis:

In environments where oxygen is absent, such as wetlands, rice paddies, sediments and anaerobic digesters, decomposition proceeds through a food chain of metabolic groups. Hydrolytic and fermentative bacteria convert polymers into sugars, amino acids and long-chain fatty acids, and then into volatile fatty acids, alcohols, carbon dioxide and hydrogen. The pattern of fermentation products depends on hydrogen partial pressure and pH, and the fermentative pathways include mixed-acid, butyrate, propionate and lactate fermentations.Syntrophic acetogenic bacteria oxidize propionate, butyrate and other intermediates to acetate, carbon dioxide and hydrogen. These reactions are endergonic under standard conditions and become feasible

only through interspecies electron transfer, in which methanogens or other hydrogen consumers keep the hydrogen and formate concentrations low. Homoacetogens add flexibility by forming acetate from hydrogen and carbon dioxide through the Wood-Ljungdahl pathway, which also functions in the oxidation of acetate in some syntrophic partnerships.The terminal step is methanogenesis carried out by archaea. Acetoclastic methanogens split acetate to methane and carbon dioxide, hydrogenotrophic methanogens reduce carbon dioxide with hydrogen, and methylotrophic methanogens use methylated compounds. Unique coenzymes including methanofuran, tetrahydromethanopterin, coenzyme M and coenzyme F430 participate in these reactions, and the final step catalyzed by methyl coenzyme M reductase releases methane. In the presence of alternative electron acceptors such as nitrate, ferric iron and sulfate, respiratory bacteria out-compete methanogens, so the dominant end products shift from methane to nitrogen, ferrous iron or hydrogen sulfide.

Environmental Controls, Applications and Future Directions:

The rate of biochemical decomposition is controlled by temperature, moisture, oxygen, pH, nutrient availability and the chemical quality of the substrate. Enzyme-catalyzed reactions generally accelerate with temperature up to an optimum, which explains the faster decomposition in warm and moist climates and the sensitivity of soil carbon stocks to climate warming. Water is required for enzyme diffusion and substrate transport, but excess water limits oxygen supply and shifts metabolism toward anaerobic pathways. Mineral surfaces and aggregates can protect organic molecules from enzymes, so that persistence of organic matter is now understood as an ecosystem property and not only a consequence of intrinsic molecular recalcitrance.Knowledge of these pathways supports many applications. In agriculture, composting and the use of microbial inoculants accelerate the conversion of crop residues into stable humus and plant-available nutrients, a practice of considerable importance for Pakistan, where large quantities of rice straw, wheat straw and sugarcane bagasse are generated each year and are often burned in the field. In energy production, anaerobic digestion converts animal manure and food waste into biogas, and enzymatic saccharification of lignocellulose provides sugars for bioethanol and other chemicals. In environmental technology, aromatic catabolic pathways are exploited for the bioremediation of petroleum hydrocarbons, pesticides and industrial dyes.Several challenges remain. The detailed mechanisms of lignin depolymerization, especially in bacteria, are still incompletely understood, and the cost of enzymes is a major obstacle for lignocellulosic biorefineries. Metagenomics, metatranscriptomics, proteomics and metabolomics are now revealing uncultured organisms and new enzyme families that participate in decomposition, and protein engineering and synthetic biology offer routes to more efficient and robust catalysts. Combining these molecular tools with ecosystem-scale measurements will be necessary to predict how decomposition and associated greenhouse gas emissions respond to land-use and climate change.



Summary:

The decomposition of complex organic substances is a multistage biochemical process in which extracellular hydrolases and oxidases depolymerize cellulose, hemicellulose, lignin, proteins and lipids, and intracellular pathways such as glycolysis, beta-oxidation, aromatic ring cleavage and the tricarboxylic acid cycle convert the products to energy, biomass and simple end products. In the absence of oxygen, fermentative, syntrophic and methanogenic communities carry out the same task through cooperative metabolic chains. The rate and direction of these pathways depend on substrate chemistry, microbial community composition and environmental conditions. Better understanding of the enzymes and regulatory networks involved will improve composting, biogas and biofuel production, bioremediation and predictions of the carbon cycle under a changing climate.

References:

- Lynd, L. R., Weimer, P. J., van Zyl, W. H., and Pretorius, I. S. (2002). Microbial cellulose utilization: fundamentals and biotechnology. Microbiology and Molecular Biology Reviews, 66, 506-577.
- Bugg, T. D. H., Ahmad, M., Hardiman, E. M., and Rahmanpour, R. (2011). Pathways for degradation of lignin in bacteria and fungi. Natural Product Reports, 28, 1883-1896.
- Cragg, S. M., Beckham, G. T., Bruce, N. C., Bugg, T. D. H., et al. (2015). Lignocellulose degradation mechanisms across the Tree of Life. Current Opinion in Chemical Biology, 29, 108-119.
- Berg, B., and McClaugherty, C. (2008). Plant Litter: Decomposition, Humus Formation, Carbon Sequestration, 2nd edition. Springer, Berlin.
- Schimel, J. P., and Bennett, J. (2004). Nitrogen mineralization: challenges of a changing paradigm. Ecology, 85, 591-602.
- Harwood, C. S., and Parales, R. E. (1996). The beta-ketoadipate pathway and the biology of self-identity. Annual Review of Microbiology, 50, 553-590.
- Thauer, R. K., Kaster, A.-K., Seedorf, H., Buckel, W., and Hedderich, R. (2008). Methanogenic archaea: ecologically relevant differences in energy conservation. Nature Reviews Microbiology, 6, 579-591.
- Cantarel, B. L., Coutinho, P. M., Rancurel, C., Bernard, T., Lombard, V., and Henrissat, B. (2009). The Carbohydrate-Active EnZymes database (CAZy): an expert resource for glycogenomics. Nucleic Acids Research, 37, D233-D238.
- Sinsabaugh, R. L. (1994). Enzymic analysis of microbial

pattern and process. Biology and Fertility of Soils, 17, 69-74.

- Lehmann, J., and Kleber, M. (2015). The contentious nature of soil organic matter. Nature, 528, 60-68.
- Kirk, T. K., and Farrell, R. L. (1987). Enzymatic combustion: the microbial degradation of lignin. Annual Review of Microbiology, 41, 465-505.
- Fuchs, G., Boll, M., and Heider, J. (2011). Microbial degradation of aromatic compounds: from one strategy to four. Nature Reviews Microbiology, 9, 803-816.