Open Horizon Scientific Review

Vol. 03 No.03 2025

Spectroscopic Investigation of Chemical Transformations in Complex Aqueous Systems

Sana Fatima Qureshi [1]

[1] Institute of Chemical Sciences, Bahauddin Zakariya University, Multan, Pakistan

ABSTRACT

Chemical transformations in natural waters, industrial effluents and biological fluids occur in mixtures containing dissolved salts, organic matter, metal ions, suspended particles and reactive intermediates. Following these changes in real time and without disturbing the sample is a major analytical challenge. This article examines how ultraviolet-visible absorption, infrared, Raman, nuclear magnetic resonance, fluorescence and mass spectrometric methods can be combined to investigate hydrolysis, photodegradation, redox reactions and metal complexation in complex aqueous systems. The strengths and limits of each technique are discussed in relation to water absorption, overlapping signals, low concentrations and matrix effects. Chemometric tools, including multivariate curve resolution and parallel factor analysis, are presented as essential means of separating overlapping spectra and extracting kinetic information. A representative case of dye degradation illustrates how spectral decay can be converted into rate constants and mechanistic insight. The numerical values shown are illustrative and serve to explain the analysis. The article concludes that integrated multi-technique spectroscopy, supported by careful data treatment, provides the most reliable route to understanding aqueous chemical change and supports better water quality monitoring and treatment design.

Key words: *Aqueous spectroscopy, Reaction monitoring, Vibrational spectroscopy, Chemometrics, Environmental chemistry*

Introduction:

Water is the medium in which most natural and many industrial chemical processes take place. Rivers, groundwater, wastewater from textile and leather industries, and biological fluids are all complex aqueous systems in which many species coexist and react. Pollutants degrade under sunlight, metals change oxidation state, organic molecules hydrolyse, and complexes form and dissociate, often within seconds or minutes. For a country such as Pakistan, where industrial effluents and agricultural runoff place heavy pressure on water resources, understanding these transformations is closely tied to public health and environmental protection.

Spectroscopy is well suited to this task because it probes molecules directly in solution, usually with little or no sample preparation, and many methods allow continuous monitoring. Each technique responds to a different

molecular property. Electronic absorption and fluorescence report on chromophores, vibrational spectroscopy on bonds and functional groups, nuclear magnetic resonance on the local environment of nuclei, and mass spectrometry on molecular composition.

However, water itself is a challenging solvent. It absorbs strongly in the infrared, scatters light when particles are present, and its hydrogen-bond network influences every solute. Real samples add further complexity in the form of dissolved organic matter and a wide range of concentrations. This article reviews how the main spectroscopic techniques are applied to aqueous chemical change, how their data are interpreted, and how they can be combined, before presenting a case study, discussing limitations and indicating future directions.

1. Nature of Complex Aqueous Systems

A complex aqueous system is defined by the presence of many interacting components rather than by a single solute in pure water. Natural waters contain inorganic ions such as carbonate, chloride, sulphate and nitrate, dissolved organic matter made of humic and fulvic substances, trace metals and microorganisms. Industrial effluents add dyes, surfactants, heavy metals and organic solvents, often at extreme pH values and high ionic strength.

The water structure itself responds to solutes. Ions and polar molecules reorganise the hydrogen-bond network, shifting vibrational bands and changing the dynamics of solvation shells. Because chemical reactivity depends on this environment, spectroscopic signals measured in water can differ significantly from those measured in organic solvents or the gas phase.

Transformations in such systems include hydrolysis of esters and amides, photolysis by sunlight, oxidation by dissolved oxygen, chlorine or hydroxyl radicals, reduction of metals, and formation of complexes with organic ligands. These processes often happen in parallel and produce transient intermediates, so a useful method must be selective, sensitive and fast enough to follow them without altering the sample.

2. Ultraviolet and Visible Absorption Spectroscopy for Reaction Monitoring

Ultraviolet and visible absorption spectroscopy is the most widely used technique for monitoring reactions in water because instruments are inexpensive, robust and compatible with flow cells and fibre-optic probes. Absorbance is linked to concentration through the Beer-Lambert law, so decay of a reactant band or growth of a product band can be turned into a kinetic curve.

In complex waters the technique provides more than simple

concentrations. Specific ultraviolet absorbance, the absorbance at 254 nanometres normalised to organic carbon, indicates the aromaticity and reactivity of dissolved organic matter, while spectral slope ratios track molecular weight and the extent of photobleaching. Metal complexation often shifts or creates charge transfer bands, allowing stoichiometry and stability constants to be determined from titration series.

The main limitation is spectral overlap. Many species absorb broadly in the ultraviolet region, and turbidity adds a sloping baseline from scattering. Derivative spectra, baseline correction, filtration and the use of full spectra rather than single wavelengths help, and diode array detectors allow entire spectra to be collected within milliseconds, which supports multivariate analysis.

3. Infrared and Raman Vibrational Spectroscopy

Vibrational spectroscopy identifies functional groups and follows bond changes directly, making it valuable for mechanistic studies. Infrared absorption by liquid water is extremely strong, especially near the bending mode at about 1640 reciprocal centimetres and the stretching region near 3400, so transmission cells need very short path lengths. Attenuated total reflectance accessories overcome this by probing only a thin layer next to a crystal and are now standard for aqueous infrared work.

Raman spectroscopy is naturally well suited to water because water is a weak Raman scatterer. Bands from solutes such as carbonate, sulphate, nitrate and organic acids can be observed clearly, and fibre-optic probes permit in situ monitoring in reactors and pipelines. Weak signals and fluorescence background are the main difficulties, and these can be reduced by using near-infrared excitation or enhanced by surface-enhanced Raman substrates, which detect trace pollutants at very low concentration.

The two methods are complementary because they follow different selection rules. Polar bonds such as carbonyl and hydroxyl groups give strong infrared bands, while symmetric vibrations and carbon-carbon bonds are strong in Raman spectra. Together they provide a fuller picture of how functional groups form and disappear during hydrolysis, oxidation or polymerisation. Studies of the water band itself also reveal changes in hydrogen bonding that accompany ion hydration and temperature shifts.

4. Nuclear Magnetic Resonance Studies of Solution Chemistry

Nuclear magnetic resonance offers detailed structural information and is inherently quantitative, since signal area is proportional to the number of nuclei.

Proton, carbon, phosphorus and fluorine spectra can identify reactants, products and intermediates in a single experiment and can track their concentrations over time without calibration curves.

The main practical difficulty in water is the huge solvent signal, which is suppressed using presaturation or selective pulse sequences, often with deuterated water added for field locking. Sensitivity is lower than for optical methods, so the technique suits millimolar rather than trace concentrations, though preconcentration or cryogenic probes can extend its reach. Paramagnetic metals broaden lines and complicate spectra, which can in turn be used to study metal binding.

Advanced experiments extend the value of the method. Diffusion-ordered spectroscopy separates species by size, which helps to distinguish free and bound molecules in mixtures, and two-dimensional correlation experiments resolve crowded spectra of natural organic matter. Rapid mixing and flow nuclear magnetic resonance setups allow reactions with half-lives of seconds to be followed in real time.

5. Fluorescence and Time-Resolved Techniques

Fluorescence spectroscopy is highly sensitive, often reaching nanomolar concentrations, and is therefore ideal for dilute natural waters. Excitation-emission matrices, which record emission over a range of excitation wavelengths, give a fingerprint of fluorescent components and allow protein-like, humic-like and fulvic-like fractions of dissolved organic matter to be distinguished.

Changes in the fluorescence landscape reveal transformations such as photobleaching, microbial processing and oxidation during water treatment. Quenching of fluorescence by metals or other solutes provides a route to binding constants, and shifts in emission wavelength report on the polarity of the local environment. Fluorescent probes designed for specific ions, pH or reactive oxygen species give selective sensing in complex matrices.

Time-resolved methods add a temporal dimension. Fluorescence lifetimes distinguish static from dynamic quenching, and transient absorption spectroscopy, using ultrafast or nanosecond laser pulses, detects short-lived intermediates such as excited states, radicals and solvated electrons that govern photochemical pathways. Inner filter effects at higher absorbance must be corrected to avoid distorted intensities.

6. Mass Spectrometric and Complementary Methods

Mass spectrometry is not strictly a spectroscopic method in the optical sense, but it is indispensable for identifying transformation products.

Electrospray ionisation allows polar and ionic species to be transferred from water into the gas phase with little fragmentation, and high-resolution instruments assign molecular formulae to hundreds or thousands of components in dissolved organic matter and degradation mixtures.

Coupling liquid chromatography with mass spectrometry separates isomers and reduces matrix suppression, while tandem mass spectrometry provides fragmentation patterns that support structural assignment. Direct online sampling interfaces allow reactions to be tracked within seconds, linking optical changes to specific molecular events.

Other complementary techniques include electron paramagnetic resonance, which detects radicals directly, X-ray absorption spectroscopy, which determines oxidation state and coordination of metals in solution, and electrochemical methods that follow redox transformations. The strongest studies combine at least two or three of these approaches so that each compensates for the weaknesses of the others.

7. Data Analysis and Chemometric Treatment of Overlapping Spectra

Complex aqueous spectra rarely contain isolated bands, so data analysis is as important as measurement. Preprocessing steps include smoothing with Savitzky-Golay filters, baseline correction, scatter correction, normalisation and subtraction of the water contribution, and each step must be applied consistently to avoid introducing artefacts.

Multivariate methods then separate overlapping contributions. Principal component analysis reveals how many independent species contribute to a data set, and multivariate curve resolution with alternating least squares extracts pure spectra and concentration profiles using constraints such as non-negativity. For fluorescence matrices, parallel factor analysis decomposes the data into components that are chemically meaningful, provided the data have the right structure.

Kinetic information is obtained by fitting concentration profiles to rate laws or by applying global analysis to the entire data set at once. Model selection must be justified, because several mechanisms may fit the same curve, and validation with independent measurements and replicate experiments gives confidence that the extracted parameters reflect real chemistry and not mathematical convenience.

8. Case Study of Dye Degradation and Other Transformations

Textile effluents provide a practical example. In a representative experiment, a coloured dye in water is exposed to ultraviolet light with and

without a semiconductor photocatalyst, and a diode array spectrometer records the visible band every few minutes. A dark control confirms that the dye is stable without light, and the decay of the main band under irradiation shows the photolysis rate. The figure below compares normalised concentration over time for the three conditions.

With light alone the colour fades slowly, whereas with the photocatalyst it disappears much faster, and fitting the curves to a first-order law gives apparent rate constants that differ by roughly an order of magnitude in the illustrative data. Disappearance of colour, however, does not prove mineralisation. Infrared and Raman spectra show loss of the aromatic and azo bands and growth of carboxylate features, while liquid chromatography coupled with mass spectrometry identifies smaller aromatic and aliphatic intermediates. Fluorescence and absorbance of the organic residue reveal whether coloured by-products form and persist.

Similar combined strategies apply to other transformations. Hydrolysis of pesticides is followed by nuclear magnetic resonance and mass spectrometry, reduction of chromium from the toxic hexavalent to the less toxic trivalent form by absorption of the diphenylcarbazide complex and X-ray absorption, and complexation of copper by humic substances by fluorescence quenching titration. In each case, no single method tells the whole story, and the integration of techniques is what gives a reliable mechanism.


9. Challenges, Environmental Applications and Future Directions

Several challenges remain. Detection limits of many techniques are above the concentrations of trace contaminants in natural waters, matrix effects complicate calibration, and the heterogeneity of dissolved organic matter makes assignment of components uncertain. Sampling and storage can also change the chemistry, which argues for in situ and online measurements wherever possible.

The applications are broad. Spectroscopic sensors can monitor drinking water quality, track the progress of advanced oxidation and disinfection, detect leaks and illegal discharges in industrial effluents, and follow nutrient and carbon cycling in rivers and lakes. Low-cost portable spectrometers and smartphone-based readers offer particular promise for regions that lack extensive laboratory infrastructure.

Future progress is likely to come from several directions: miniaturised and fibre-based sensors, ultrafast and two-dimensional spectroscopies that reveal solvation dynamics, enhanced signals from nanostructured substrates, and machine learning tools that classify spectra and predict composition from

large data sets. Close integration of experiment, computation and standardised data treatment will make results more comparable between laboratories and more useful for regulators and water managers.



Summary:

This article reviewed the spectroscopic investigation of chemical transformations in complex aqueous systems. Ultraviolet-visible absorption offers accessible kinetic monitoring, vibrational methods reveal bond-level changes, nuclear magnetic resonance provides quantitative structural detail, fluorescence gives high sensitivity, and mass spectrometry identifies products. Water absorption, spectral overlap, low concentration and matrix effects are the main obstacles, and chemometric methods such as multivariate curve resolution and parallel factor analysis are essential to overcome them. A dye degradation example showed how kinetic curves from several conditions can be compared and how supporting techniques confirm the mechanism. Combining several spectroscopic methods with careful data treatment gives the most dependable understanding of aqueous chemical change and supports monitoring, treatment and protection of water resources.

References:

- Bakker, H. J., & Skinner, J. L. (2010). Vibrational spectroscopy as a probe of structure and dynamics in liquid water. Chemical Reviews, 110, 1498–1517.
- Coble, P. G. (1996). Characterization of marine and terrestrial DOM in seawater using excitation-emission matrix spectroscopy. Marine Chemistry, 51, 325–346.
- de Juan, A., & Tauler, R. (2006). Multivariate curve resolution (MCR) from 2000: progress in concepts and applications. Critical Reviews in Analytical Chemistry, 36, 163–176.
- Eisenberg, D., & Kauzmann, W. (1969). The Structure and Properties of Water. Oxford University Press, Oxford.
- Günther, H. (2013). NMR Spectroscopy: Basic Principles, Concepts and Applications in Chemistry (3rd ed.). Wiley-VCH, Weinheim.
- Helms, J. R., Stubbins, A., Ritchie, J. D., Minor, E. C., Kieber, D. J., & Mopper, K. (2008). Absorption spectral slopes and slope ratios as

indicators of molecular weight, source, and photobleaching of chromophoric dissolved organic matter. Limnology and Oceanography, 53, 955–969.

- Lakowicz, J. R. (2006). Principles of Fluorescence Spectroscopy (3rd ed.). Springer, New York.
- Savitzky, A., & Golay, M. J. E. (1964). Smoothing and differentiation of data by simplified least squares procedures. Analytical Chemistry, 36, 1627–1639.
- Skoog, D. A., Holler, F. J., & Crouch, S. R. (2017). Principles of Instrumental Analysis (7th ed.). Cengage Learning, Boston.
- Smith, E., & Dent, G. (2019). Modern Raman Spectroscopy: A Practical Approach (2nd ed.). Wiley, Chichester.
- Stedmon, C. A., & Bro, R. (2008). Characterizing dissolved organic matter fluorescence with parallel factor analysis: a tutorial. Limnology and Oceanography: Methods, 6, 572–579.
- Weishaar, J. L., Aiken, G. R., Bergamaschi, B. A., Fram, M. S., Fujii, R., & Mopper, K. (2003). Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon. Environmental Science and Technology, 37, 4702–4708.