Open Horizon Scientific Review

Vol. 04 No.02 2026

Health Information Exchange and Continuity of Care Across Healthcare Institutions

Saad Hamza Qureshi

Department of Public Health, Health Services Academy (HSA), Islamabad, Pakistan.

ABSTRACT

Continuity of care becomes difficult when patients receive services from multiple healthcare institutions that maintain separate clinical records. Important information may arrive late, remain inaccessible, or lack the context necessary for safe interpretation. Health information exchange provides an organizational and technical means of sharing clinical information across institutional boundaries, but data transmission alone does not ensure coordinated care. This narrative review examines the relationship between health information exchange and informational, management, and relational continuity. Selected scholarly literature, an interoperability specification, and international digital health guidance inform an analytical discussion of exchange architecture, data quality, patient identification, transitions of care, clinical workflow, privacy, implementation, and evaluation. The article distinguishes technical connectivity from meaningful clinical use and separates process improvements from demonstrated patient benefit. A conceptual framework proposes that successful exchange requires a sequence of reliable patient identification, relevant information availability, clinician interpretation, assigned responsibility, completed action, and patient understanding. Particular attention is given to the planning considerations relevant to Pakistani healthcare institutions, without claiming nationally representative findings or reporting an original patient dataset. A graph presents the distribution of study designs in a published systematic review and illustrates limitations in the evidence base rather than an intervention effect. The discussion identifies practical priorities including accurate medication information, timely discharge communication, referral tracking, correction of shared errors, equitable access, and sustainable funding. Health information exchange can strengthen continuity when embedded within accountable care processes, but effectiveness depends on local implementation, trustworthy governance, and evaluation of outcomes beyond the volume of exchanged records.

Key words : *Health information exchange, Continuity of care, Interoperability, Care coordination, Patient safety*

Introduction:

Healthcare frequently involves a series of encounters across general practices, emergency departments, hospitals, laboratories, pharmacies, rehabilitation services, and specialist clinics. Each institution may possess only part of the information required to understand the patient's condition and treatment history. When these fragments remain disconnected, clinicians may repeat investigations, overlook

medication changes, misunderstand previous decisions, or fail to complete follow-up. Patients and families may be expected to reconstruct the missing history while dealing with illness, financial pressures, and uncertainty. The resulting discontinuity is an organizational problem as well as an information problem because information must reach someone who can interpret it and act.

Health information exchange refers to the electronic sharing of health information across organizational boundaries. The term can describe an activity, a technical capability, or an arrangement through which institutions agree to share information. These meanings overlap but should not be treated as identical. A hospital may possess an interface capable of sending documents while lacking a dependable process for identifying recipients and confirming follow-up. Conversely, a small referral network may achieve useful continuity through a limited but carefully managed exchange service. The appropriate unit of analysis is therefore the complete care process supported by exchange, rather than the network connection alone.

Continuity of care provides a useful framework for examining this process. Haggerty et al. (2003) distinguish informational, management, and relational continuity, emphasizing that continuity is experienced over time. Shared information can support all three dimensions, but it cannot replace a coherent care plan or a trusted professional relationship. A complete record may still leave patients uncertain about who is responsible for their next appointment. Equally, strong relationships may be undermined when clinicians cannot access important information from another institution. Effective exchange should therefore connect clinical knowledge with responsibility, communication, and the patient's experience of coordinated care.

This article is a narrative review and conceptual analysis based on selected publications relevant to exchange, continuity, transitions, interoperability, and implementation. It does not report exhaustive searching, a registered protocol, formal study selection, or a new meta-analysis. Published reviews provide an evidence foundation, while the proposed implementation framework and examples represent analytical synthesis. The central question is how information exchange can contribute to safer and more coherent care across institutions, particularly when resources and digital capabilities differ. Attention is given to Pakistani implementation considerations as proposals for local assessment rather than as conclusions from a national survey.

1 Conceptual Relationship Between Information Exchange and Continuity

Informational continuity means that knowledge from previous encounters contributes meaningfully to current care. This includes observations, test results, medication history, patient preferences, and the reasoning behind earlier decisions. A document that is available but difficult to locate may provide little practical continuity. Similarly, a result without its date, units, or clinical context may be interpreted incorrectly. Exchange should therefore preserve meaning and relevance as well as content. The clinical objective is to reduce the need to reconstruct essential history while maintaining awareness that previously recorded information may be incomplete, uncertain, or no longer applicable.

Management continuity concerns a coherent approach across professionals and settings. A patient discharged from hospital may need medication review, laboratory monitoring, rehabilitation, and specialist follow-up. These actions form a connected plan only when responsibility, timing, and communication are explicit. Information exchange can make the plan visible, but visibility does not establish acceptance of responsibility. Receiving institutions may require a process for confirming that a referral has been reviewed and an appointment arranged. A useful framework separates the stages of sending, receiving, understanding, accepting, completing, and communicating an action. Failures at any stage can interrupt continuity despite successful electronic transmission.

Relational continuity involves an ongoing therapeutic relationship with a clinician or team. Exchange can support this relationship by allowing a familiar professional to remain informed during treatment elsewhere. It can also reduce the burden of repeatedly describing distressing experiences. However, indiscriminate sharing may weaken trust when patients do not understand who can access their information. A patient might prefer a sensitive concern to be discussed directly rather than encountered unexpectedly in a broad record. Relational continuity therefore requires transparent communication about sharing and a response to patient concerns. Technical availability should strengthen trust through appropriate use and accountable stewardship.

The conceptual model proposed here links six necessary conditions: accurate identification, relevant information availability, understandable presentation, accountable interpretation, completed clinical action, and patient understanding. These conditions are not a validated scoring instrument. They are a practical way to identify where an

exchange initiative may fail to influence care. An institution could achieve excellent network uptime while struggling with incorrect patient matches or unassigned follow-up tasks. Measuring each condition separately can prevent technical success from concealing clinical failure. It also supports targeted improvement because organizations can address a specific weak point instead of repeatedly investing in broader connectivity.

2 Exchange Architecture and Technical Interoperability

Exchange arrangements differ in how information is stored, discovered, and delivered. A centralized arrangement may maintain shared copies of selected information, while a federated arrangement may retrieve information from participating institutions when requested. Hybrid arrangements combine elements of both. Directed exchange sends information to an intended recipient, whereas query-based exchange supports searching for relevant external records. Event notifications may inform a care team that a patient has been admitted or discharged. These approaches serve different purposes and create different operational demands. Architecture should therefore be selected around a defined care pathway, expected information needs, and institutional responsibilities. Vest and Gamm (2010) similarly emphasize challenges extending beyond technology.

Technical interoperability involves the ability of systems to communicate, while semantic interoperability concerns whether the exchanged information retains a shared meaning. Common message structures and interfaces help systems transfer data, but agreement on codes, units, status, and context remains necessary. A laboratory result represented as text may be readable by a clinician but unsuitable for automated comparison. Conversely, a structured value can be misleading if its unit or specimen context is omitted. Institutions should agree on a minimum clinically meaningful dataset and test how receiving systems display it. A technically accepted message is not sufficient evidence of clinically correct interpretation.

The HL7 FHIR specification provides a resource-based framework for representing and exchanging healthcare information. Reference to FHIR does not imply that every implementation supports the same resources, operations, terminology bindings, or clinical workflows. Participating institutions still need agreed implementation details and conformance testing. D'Amore et al. (2014), examining a different document-based exchange framework, demonstrated why nominal standards compliance should not be equated with

reliable interoperability. The broader lesson is that standards need practical testing against realistic records. A procurement requirement should therefore ask what information can be exchanged correctly in the intended setting, rather than merely whether a standard is named.

Reliability also requires clear behavior when exchange is unavailable or incomplete. A clinician should know whether no external records exist, a query failed, access was restricted, or an institution is temporarily offline. Treating all these situations as an empty history can create false reassurance. Systems should preserve the source and retrieval time of information and avoid implying that a partial record represents the whole patient journey. Downtime procedures should specify how urgent communication proceeds and how delayed information is reconciled afterward. Architecture becomes clinically useful when it communicates its own limits and supports dependable recovery from routine failures.

3 Patient Identification, Data Quality, and Shared Meaning

Patient identification is foundational because accurate information becomes harmful when attached to the wrong person. Matching may rely on combinations of demographic details, local identifiers, and other approved attributes. Institutions must examine the risks of both false matches and missed matches. A false match can introduce another person's diagnoses or medications, while a missed match can hide important history. These errors have different consequences and require different responses. Matching performance should be evaluated with representative local data, including variations in spelling, incomplete fields, duplicate records, and changes in contact details. Confidence should not be inferred from a successful technical connection.

Record linkage should include human review for uncertain cases and a clear method for reversing incorrect associations. Staff need to understand which details can be corrected locally and how corrections propagate across institutions. A mistaken merge that is fixed in one system may remain visible elsewhere unless there is a coordinated correction process. Audit trails should identify what changed, why it changed, and which organizations received the correction. These processes require assigned responsibility and accessible support. Identity management is therefore an ongoing service rather than a one-time registration exercise, especially when patients move between facilities with different recordkeeping practices.

Data quality extends beyond correct values. Completeness, timeliness,

provenance, internal consistency, and clinical relevance determine whether information supports continuity. A medication list should distinguish active therapy from discontinued or historical prescriptions. An allergy record should separate a documented reaction from an unverified report. A diagnosis should indicate whether it is confirmed, suspected, resolved, or copied from an earlier encounter. Without these distinctions, exchange can amplify ambiguity by making uncertain information appear authoritative. Systems should display uncertainty and origin clearly, allowing clinicians to examine the basis of a shared statement before incorporating it into current decisions.

Shared meaning also depends on preservation of narrative context. Structured fields can support searching and comparison, while concise clinical narratives explain why a decision was made. Neither format is sufficient in every circumstance. A discharge diagnosis may be less useful without the unresolved questions and reasons for medication changes. Institutions should avoid reducing continuity to an accumulation of coded observations. A balanced exchange record combines structured essentials with readable explanations and clear source attribution. Quality monitoring should examine whether clinicians can understand and use the information, not simply whether mandatory fields contain some text or a code.

4 Transitions, Referrals, and Longitudinal Care Coordination

Transitions between institutions are particularly vulnerable because information, professional responsibility, and the patient's location change at the same time. Kripalani et al. (2007) reviewed deficits in communication between hospital-based and primary care physicians, identifying the importance of discharge information for continuity. The relevance of this work is the enduring need for timely and usable handover. A contemporary exchange service should make essential information available when the receiving team needs it, while avoiding unsupported assumptions that a historical study describes current practice in every healthcare system. Local baseline assessment remains necessary before choosing an intervention or claiming improvement.

A useful discharge communication includes the reason for admission, important findings, treatment changes, pending results, unresolved problems, and a clear follow-up plan. The receiving clinician also needs to understand which actions are urgent and who has agreed to undertake them. If a test remains pending, the plan should identify the team responsible for

reviewing the result and communicating with the patient. Merely placing the result in a shared repository does not close the loop. An effective exchange pathway should therefore connect documents with worklists or other operational processes that track acceptance and completion without creating duplicate or contradictory assignments.

Referral exchange should support communication in both directions. The referring clinician needs to know whether a referral was received, triaged, scheduled, and completed, while the specialist needs enough information to assess the reason for consultation. After the visit, the specialist's conclusions should return to the team responsible for ongoing care. These stages are especially important when patients attend a different institution from the one originally intended. A network should establish how external outcomes are captured and how unresolved referrals are reviewed. The objective is a coherent patient journey rather than a count of referral messages successfully transmitted between servers.

Care coordination also includes patients and caregivers. Coleman et al. (2006) evaluated a care transitions intervention involving patient participation and coaching; that trial should not be interpreted as an isolated test of electronic exchange. Its relevance is that continuity interventions often depend on human support as well as information tools. A shared record can assist a transition coach, nurse, or general practitioner, but it does not replace explanation and practical assistance. Patients should leave an encounter understanding the next step and the contact route for problems. Support should accommodate those who cannot use a portal or manage multiple appointments independently.

5 Evidence of Clinical Benefit and Limits of Existing Research

Published reviews provide reasons for cautious optimism while showing why broad claims require qualification. Rudin et al. (2014) reviewed the usage and effects of exchange, and Hersh et al. (2015) examined outcomes and research needs. Rahurkar, Vest, and Menachemi (2015) highlighted limitations in evidence on cost, utilization, and quality. These publications address heterogeneous exchange arrangements and clinical settings. Their findings should not be combined informally into a single expected percentage improvement in continuity. Availability of records, frequency of consultation, reduction in duplicated tests, and improvement in patient health are distinct outcomes with different evidential requirements.

Menachemi et al. (2018) subsequently reported a more favorable

picture among stronger study designs in an updated review. Their review included 24 articles, of which seven used designs considered suitable for causal inference; those seven reported some beneficial effect. This is not equivalent to proof that all exchange services improve outcomes or that seven randomized trials demonstrated a universal benefit. The types of exchange, implementation settings, outcomes, and risks of bias remain relevant. A balanced interpretation recognizes positive evidence while asking whether the local service resembles the interventions studied and whether the intended outcome was actually measured.

The accompanying graph shows seven studies classified by that review as suitable for causal inference and seventeen other included studies, together accounting for the twenty-four articles. The second value is calculated by subtraction. The graph describes the composition of the review, not the magnitude of an intervention effect, a success rate, or the proportion of patients helped. It is included to illustrate why study design matters when interpreting an evidence base. Readers should not treat the two bars as clinical comparison groups. No patient-level values, confidence intervals, or before-and-after performance estimates have been invented for the figure.

6 Workflow, Professional Roles, and Patient Participation

Clinicians are more likely to use shared information when it is accessible within the task they are performing. Separate logins, unfamiliar interfaces, long document lists, and slow retrieval can make technically available information practically inaccessible. Eden et al. (2016) identified workflow and early user involvement among relevant factors in exchange use. For implementation, the practical implication is to observe real clinical work before finalizing an interface. The aim is to place relevant information near the decision it supports, with enough context to be understood and enough restraint to avoid overwhelming the user with unrelated historical material.

Information presentation should prioritize clinical relevance without concealing the source. A concise summary may highlight recent admissions, medication changes, important results, and unresolved actions. Users should be able to inspect the underlying document when needed. Automated summarization requires validation because omission or misinterpretation can distort the record. Even a manually created summary becomes outdated unless there is a process for updating it. The interface should therefore distinguish source information from derived summaries and

show their dates. Searchability and readability should be tested using realistic cases, including conflicting records and patients with lengthy histories across several institutions.

Professional responsibilities must be agreed across organizational boundaries. A receiving team may have access to a result without having accepted responsibility for follow-up. Shared access should not create an assumption that someone else has acted. Institutional agreements should identify how tasks are assigned, when they are acknowledged, and what happens if the recipient cannot respond. Staff need protected time and clear escalation routes. The exchange service should support these arrangements but cannot substitute for them. Evaluation should examine whether the distribution of work is sustainable and whether information exchange has silently increased administrative burdens for particular teams.

Patient participation can improve the accuracy and usefulness of shared records. Patients may identify an outdated medication, clarify an encounter at another facility, or report that a referral did not occur. Their contributions should enter a review process that distinguishes reported information from verified clinical updates. Access arrangements should accommodate disability, language, literacy, and availability of digital devices. A portal should be one route for participation rather than the only route. Patients should also receive explanations of discrepancies when they affect care. Participation is meaningful when it helps resolve a problem and produces a response, not merely when the system provides a feedback button.

7 Privacy, Trust, Security, and Institutional Governance

Trust is central to exchange because institutions are sharing information that may have been collected in a different clinical relationship. Patients should understand the purposes of sharing and the categories of professionals who may access their information. Consent and authorization arrangements depend on applicable law and policy and require local legal review. This article does not prescribe a universal legal model for Pakistan or other jurisdictions. Instead, it recommends clear documentation of the chosen arrangements, practical communication with patients, and processes for responding to restrictions, questions, and complaints. Ambiguous governance can undermine both participation and appropriate clinical use.

Access controls should reflect professional role, legitimate purpose, and the sensitivity of the information. Organizations need procedures for

granting, reviewing, and revoking access, especially when staff change roles or leave an institution. Emergency access, where permitted, should be limited and auditable. Security also includes secure transmission, protected storage, reliable authentication, monitoring, and incident response. These controls should be assessed as part of routine operations rather than a checklist completed only at launch. A network can be secure against external intrusion yet still expose information through excessive internal permissions or casual sharing of user accounts.

Governance must address responsibility for incorrect information and cross-institutional incidents. Participants should know who investigates a suspected mismatch, how an erroneous result is corrected, and how affected users are notified. Agreements should also specify the permitted uses of data and distinguish direct care from research, analytics, or commercial activity. Secondary uses require their own justification and safeguards. Institutional participation in exchange should not be interpreted as unrestricted permission for all future purposes. A transparent governance structure includes clinical, technical, organizational, and patient perspectives so that decisions are not determined solely by the priorities of a platform vendor or one large hospital.

The World Health Organization's Global Strategy on Digital Health 2020–2027 emphasizes a broader strategic approach to digital health rather than isolated technology acquisition. In the framework proposed here, governance should connect exchange investment with health service priorities and accountable leadership. Independent oversight may be appropriate for disputed access, safety incidents, and changes in data use. Institutions also need a clear exit process that protects continuity if a participant withdraws or a provider changes. Trust depends on predictable behavior throughout the service lifecycle, including failure, correction, transition, and closure, rather than assurances made only during procurement.

8 Implementation Priorities for Pakistani Healthcare Institutions

A proposed implementation in Pakistan should begin with a defined patient pathway and a documented baseline problem. Examples might include hospital discharge to primary care, transfer between a district facility and a specialist center, or communication of laboratory findings across a referral network. These are illustrative use cases, not claims about national prevalence. Local assessment should examine institutional workflows, existing records, staffing, language needs, infrastructure, and patient access.

A limited pathway can make responsibilities and outcomes easier to evaluate. Starting with broad data aggregation without a clear clinical purpose may create considerable technical work while leaving the continuity problem unchanged.

The minimum exchange dataset should be negotiated with intended users. Essential content might include verified identifying details, encounter information, current medications with status, relevant allergies, key findings, pending actions, and the responsible team. The list should remain proportionate to the chosen use case. Requiring every participating institution to supply a comprehensive record at the outset may exclude smaller facilities or encourage low-quality completion. A phased approach can establish a dependable core before expanding. Each phase should specify what information is available, what remains outside the network, and how clinicians should manage gaps in the shared record.

Infrastructure planning should include electricity, connectivity, device availability, technical support, and recovery procedures where these are relevant to the participating sites. These factors should be measured locally rather than assumed to be uniform across Pakistan. A system should handle delayed synchronization safely and make the age of information visible. Institutions need a workable alternative when exchange is unavailable, especially for urgent transfers. Training should cover clinical interpretation, identity verification, privacy, correction of errors, and follow-up responsibilities. Competence should be demonstrated through practical scenarios rather than inferred from attendance at a software demonstration or completion of a short presentation.

Financing should consider the costs borne by all participants. A large institution may receive substantial benefits while a smaller clinic carries additional data entry or follow-up work. Sustainable arrangements should therefore consider shared funding, support for participation, and allocation of operational responsibilities. Cost estimates should include implementation, maintenance, terminology updates, staff time, cybersecurity, training, and eventual migration. Patient expenses also matter when new workflows require additional travel or digital access. An implementation is more credible when it shows how the service will be maintained after initial funding ends and how its benefits will be assessed across institutions of different sizes.

9 Evaluation, Sustainability, and Future Development


Evaluation should begin with a theory of change that links a specific exchange function to a care outcome. For example, timely discharge information may support medication reconciliation, which may reduce clinically important discrepancies and subsequent harm. Each link should be measured where feasible. Technical measures include successful transmission, retrieval time, and data completeness. Process measures include timely review, referral completion, and resolution of pending actions. Patient measures include understanding, experience of coordination, safety, and relevant clinical outcomes. An evaluation should avoid using a high message count as a substitute for evidence that information improved care.

Study design should account for differences between institutions and changes over time. Cluster trials, controlled implementation studies, and carefully designed interrupted time-series analyses may be appropriate depending on feasibility and context. The protocol should define the intervention, comparator, primary outcome, and follow-up period in advance. Investigators should document actual use, missing data, concurrent initiatives, and unintended consequences. A network that has been installed but rarely consulted represents a different exposure from one integrated into routine care. Reporting should therefore distinguish availability, use, and completed action rather than combining them under a single binary label of exchange adoption.

Equity and sustainability should be assessed alongside effectiveness. Overall improvements can conceal weaker outcomes for patients with incomplete identifiers, limited literacy, uncommon languages, or care outside participating institutions. Evaluation should examine who is missing from the network and whether clinicians wrongly interpret those gaps as absence of prior care. Sustainability measures should include staffing, user burden, financial viability, maintenance capacity, and the ability to change vendors without losing essential information. Continuing investment should depend on demonstrated value and a practical improvement process. A system should be simplified or redesigned when complexity increases faster than useful clinical benefit.

Future development may include richer event notifications, patient contributions, structured care plans, and carefully validated decision support based on shared records. These capabilities should build on trustworthy data and clear accountability. Adding predictive analytics to incomplete or incorrectly linked records can amplify existing problems. Institutions should

first ensure that the exchange reliably supports fundamental tasks such as identifying the right patient and communicating the next step. The long-term goal is a learning network in which cross-institutional information helps teams recognize problems, coordinate action, and improve service design. Technological sophistication is valuable when it strengthens that goal and remains understandable to its users.



Source: Menachemi et al. (2018), n = 24 articles. Other studies: 24 − 7 = 17. Describes review composition; does not measure a clinical treatment effect.

Summary

Health information exchange can support continuity by making relevant clinical information available across institutional boundaries. Its contribution extends beyond transferring documents because effective continuity requires interpretation, responsibility, completed action, and understandable communication with patients. Informational continuity depends on accurate, timely, contextual records; management continuity requires coherent plans and accepted responsibilities; relational continuity relies on trust and ongoing professional relationships. A successful exchange service should support these dimensions together while recognizing that access to data does not automatically create coordinated care. The proposed framework identifies the practical stages through which shared information must pass before it can influence the patient's experience or clinical outcome.

The evidence base includes studies reporting benefits as well as methodological and implementation limitations. Published reviews should be interpreted according to their clinical settings, exchange arrangements, study designs, and outcome definitions. The graph included in this article describes the design composition of one published review; it does not estimate the effectiveness of an exchange intervention. Claims of reduced duplication, lower expenditure, better safety, or improved continuity require appropriate local evaluation. Technical uptime, message counts, and document availability remain useful operational indicators, but they cannot alone establish that care has become safer or more coherent.

For Pakistani institutions, a practical strategy is to select a defined care pathway, agree a meaningful minimum dataset, establish reliable identification, assign follow-up responsibilities, and test the service

under realistic conditions. Privacy, patient participation, multilingual communication, equitable access, and sustainable funding should be addressed from the beginning. Governance must support correction of errors, incident investigation, and accountable decisions about data use. Long-term success depends on maintaining the service and adapting it to changing clinical needs. Exchange strengthens continuity when the right information reaches an appropriate professional in time to support a justified action that the patient understands and can realistically complete.

References:

- Coleman, E. A., Parry, C., Chalmers, S., & Min, S.-J. (2006). The care transitions intervention: Results of a randomized controlled trial. Archives of Internal Medicine, 166(17), 1822–1828. https://doi.org/10.1001/archinte.166.17.1822
- D'Amore, J. D., Mandel, J. C., Kreda, D. A., et al. (2014). Are Meaningful Use Stage 2 certified EHRs ready for interoperability? Findings from the SMART C-CDA Collaborative. Journal of the American Medical Informatics Association, 21(6), 1060–1068. https://doi.org/10.1136/amiajnl-2014-002883
- Eden, K. B., Totten, A. M., Kassakian, S. Z., et al. (2016). Barriers and facilitators to exchanging health information: A systematic review. International Journal of Medical Informatics, 88, 44–51. https://pubmed.ncbi.nlm.nih.gov/26878761/
- Haggerty, J. L., Reid, R. J., Freeman, G. K., Starfield, B. H., Adair, C. E., & McKendry, R. (2003). Continuity of care: A multidisciplinary review. BMJ, 327(7425), 1219–1221. https://doi.org/10.1136/bmj.327.7425.1219
- Health Level Seven International. (2019). FHIR Release 4: Overview (Version 4.0.1). https://hl7.org/fhir/R4/overview.html
- Hersh, W. R., Totten, A. M., Eden, K. B., et al. (2015). Outcomes from health information exchange: Systematic review and future research needs. JMIR Medical Informatics, 3(4), e39. https://doi.org/10.2196/medinform.5215
- Kripalani, S., LeFevre, F., Phillips, C. O., Williams, M. V., Basaviah, P., & Baker, D. W. (2007). Deficits in communication and information transfer between hospital-based and primary care physicians: Implications for patient safety and continuity of care. JAMA, 297(8), 831–841. https://doi.org/10.1001/jama.297.8.831
- Menachemi, N., Rahurkar, S., Harle, C. A., & Vest, J. R. (2018). The benefits of health information exchange: An updated systematic review. Journal of the American Medical Informatics Association, 25(9), 1259–1265. https://doi.org/10.1093/jamia/ocy035
- Rahurkar, S., Vest, J. R., & Menachemi, N. (2015). Despite the spread of health information exchange, there is little evidence of its impact on cost, use, and quality of care. Health Affairs, 34(3), 477–483.

https://doi.org/10.1377/hlthaff.2014.0729

- Rudin, R. S., Motala, A., Goldzweig, C. L., & Shekelle, P. G. (2014). Usage and effect of health information exchange: A systematic review. Annals of Internal Medicine, 161(11), 803–811. https://doi.org/10.7326/M14-0877

- Vest, J. R., & Gamm, L. D. (2010). Health information exchange: Persistent challenges and new strategies. Journal of the American Medical Informatics Association, 17(3), 288–294. https://doi.org/10.1136/jamia.2010.003673

- World Health Organization. (2025). Global strategy on digital health 2020–2027. https://www.who.int/publications/i/item/9789240116870